

Before using this product read the label carefully

**ALEXIN<sup>®</sup>**

Reg No. B3835, Act No. 36 of 1947

Namibian Reg. No. N-F 0674

Botswana Reg. No.

**FERTILIZER GROUP 2**

**A liquid organic nutrient complex containing salicylic acid.**

**ACTIVE INGREDIENTS:**

Calcium	(Ca)	26.4 g/kg	(33.3 g/l)
Magnesium	(Mg)	8.4 g/kg	(10.6 g/l)
Potassium	(K)	45 g/kg	(56.7 g/l)
Boron	(B)	2.375 g/kg	(3.0 g/l)

**CONTENTS:**

Batch No:

Date of Manufacture:

**REGISTRATION HOLDER:**

**Nulandis<sup>®</sup> (A division of AECI Limited)**

Reg. No. 1924/002590/06

P.O. Box 14418

Witfield, 1467

Tel : (011) 823 8000



**ALEXIN<sup>®</sup>** is a registered trademark of Nulandis<sup>®</sup> (A division of AECI Limited)

**DIRECTIONS FOR USE:** USE ONLY AS INDICATED

- Alexin should not be added with any sulphur or phosphite based products

**APPLICATION TABLE:**

<b>CROP</b>	<b>APPLICATION RATE</b>	<b>APPLICATION METHOD</b>
<b>APPLES, PEARS</b>	200-300 ml/100 l water to a maximum of 3 l/ha	Apply 3-4 applications from 50% flowering and repeat with 21 day intervals. <b>For use on Pears: Do not combine Alexin with other foliar feeds or adjuvants.</b>
<b>CITRUS</b>	200-300 ml/100 l water to a maximum of 3 l/ha	Apply 3 – 4 applications from 100 % petal drop and repeat with 28 day intervals.
<b>CUCURBITS (PUMPKINS, WATERMELONS, MELONS AND CUCUMBERS)</b>	500 ml /100 l water to a maximum of 4 l/ha	Apply at flowering and repeat with 21 day intervals.
<b>LETTUCE, CAULIFLOWER, CABBAGE, CHINESE CABBAGE</b>	500 ml/100 l water to a maximum of 3 l/ha	Begin applications prior to onset of head formation and repeat at 14 to 21 day intervals.
<b>PEACHES, NECTARINES, PLUMS, APRICOTS, ALMONDS, OLIVES</b>	200-300 ml/100 l water to a maximum of 3 l/ha	Apply 3-4 applications from 75% petal drop and repeat with 21 to 28 day intervals. If weather conditions favour bacterial infections, additional applications can be considered.
<b>POTATOES</b>	500 ml/100 l water to a maximum of 4 l/ha	Apply at tuber initiation and repeat every 21 to 28 days.
<b>TOMATOES, PEPPERS, BEANS, PEAS, ONIONS</b>	500 ml/100 l water to a maximum of 4 l/ha	Begin applications prior to fruit set and repeat at 14 to 21 day intervals.
<b>TOBACCO</b>	500 ml/100 l water to a maximum of 4 l/ha	Apply 2 – 3 applications bi-weekly, starting at 30 cm height.

**IMPORTANT: BEFORE APPLYING ALEXIN, CONSULT A QUALIFIED AGRONOMIST OR AGRICULTURAL ADVISER ON THE USE THEREOF.**

# ALEXIN<sup>®</sup>

Reg Nr. B3835, Wet Nr. 36 van 1947

Namibiese Reg. Nr. N-F 0674

Botswana Reg. Nr.

## 'n Vloeibare organiese voedingskompleks wat salisielsuur bevat. MISSTOF GROEP 2

### AKTIEWE BESTANDELE:

Kalsium	(Ca)	26.4 g/kg (33.3 g/l)
Magnesium	(Mg)	8.4 g/kg (10.6 g/l)
Kalium	(K)	45 g/kg (56.7 g/l)
Boor	(B)	2.375 g/kg (3.0 g/l)

INHOUD:

Lotnommer:

Datum van Vervaardiging:

### REGISTRASIEHOUER:

Nulandis<sup>®</sup> ('n Afdeling van AECI Beperk)

Reg. Nr. 1924/002590/06

Posbus 14418

Witfield, 1467

Tel : (011) 823 8000



**ALEXIN<sup>®</sup>** is 'n geregistreerde handelsmerk van Nulandis<sup>®</sup> ('n Afdeling van AECI Beperk)

### GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- **Alexin** moenie by swael- of fosfiet-gebaseerde produkte gevoeg word nie.

### TOEDIENINGSTABEL:

GEWAS	TOEDIENINGSHOEVEELHEID	TOEDIENINGSMETODE
AARTAPPELS	500 ml/100 l water tot 'n maksimum van 4 l/ha	Dien toe tydens knolvorming en herhaal elke 21 tot 28-dae.
APPELS, PERE	200 - 300 ml/100 l water tot 'n maksimum van 3 l/ha	Dien 3-4 keer toe vanaf 50 % blom en herhaal met 21-dae tussenposes. <b>Vir gebruik op Pere: Moenie Alexin met ander blaarvoedings- of byvoegmiddels meng nie.</b>
BLAARSLAAI, BLOMKOOL, KOOL, CHINESE KOOL	500 ml/100 l water tot 'n maksimum van 3 l/ha	Begin toedien vóór kopvorming en herhaal met 14 tot 21-dae tussenposes.
PAMPOENGWASSE (PAMPOENE, WAATLEMOENE, SPANSPEKKE EN KOMKOMMERS)	500 ml/100 l water tot 'n maksimum van 4 l/ha	Dien toe tydens blom en herhaal met 21-dae tussenposes.
PERSKES, PRUIME, NEKTARIENS, AMANDELS, OLYWE, APPELKOSE	200 - 300 ml/100 l water tot 'n maksimum van 3 l/ha	Dien 3-4 keer toe vanaf 75 % blomblaarval en herhaal met 21 tot 28-dae tussenposes. Indien weersomstandighede gunstig is vir bakteriese infeksies, kan addisionele behandelings oorweeg word.
SITRUS	200 - 300 ml/100 l water tot 'n maksimum van 3 l/ha	Dien 3-4 keer toe vanaf 100 % blomblaarval en herhaal met 28-dae tussenposes.
TAMATIES, SOETRISSIES, BONE, ERTJIES, UIE	500 ml/100 l water tot 'n maksimum van 4 l/ha	Begin toedien vóór vrugset en herhaal met 14 tot 21-dae tussenposes.
TABAK	500 ml/100 l water tot 'n maksimum van 4 l/ha	Dien 2 - 3 keer toe elke 14 dae. Begin toedien op 'n hoogte van 30 cm.

**BELANGRIK: RAADPLEEG 'N GEKWALIFISEERDE AGRONOOM OF LANDBOUKUNDIGE, VOORDAT ALEXIN TOEGEDIEN WORD, OOR DIE NOODSAAKLIKHEID VAN DIE GEBRUIK DAARVAN.**