

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE PRECAUTIONS

1. Handle with care. Avoid splashing and contact with eyes, nose and mouth.
2. Poisonous by swallowing, inhalation and contact with skin.
3. **WEAR FULL PERSONAL PROTECTIVE EQUIPMENT (PPE)** i.e. rubber gloves, rubber boots, overalls, hood, goggles and respirator when preparing and using with this product.
4. **KEEP APART FROM FOOD AND FOOD STUFFS.**
5. **DO NOT** eat, drink or smoke while preparing and applying this product.
6. On completion of spraying remove personal protective clothing, wash face, hands and body with soap and running water before engaging in any activities.
7. **KEEP OUT OF REACH OF CHILDREN AND UNIFORMED PERSONS.**
8. Dangerous to livestock and domestic pets. Very dangerous to aquatic organisms and may cause long-term adverse effects in the aquatic environment.
9. Store in the original container and **KEEP UNDER LOCK AND KEY.**
10. **DO NOT** contaminate drinking pools, streams, water bodies with the chemical or empty containers.
11. Prevent drift onto other crops and areas not under treatment.

SYMPTOMS OF POISONING:

Vomiting, anxiety, salivation, shivering, convulsions and skin, eye, nose and throat irritation.

FIRST AID:

Inhalation: Move victim to fresh air. If not breathing properly, give artificial respiration. Seek medical attention.

Skin contact: Remove contaminated clothing and shoes and launder. Wash thoroughly with soap and running water.

Eye contact: Flush immediately with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

Ingestion: If swallowed, wash the mouth with water provided person is conscious. Induce vomiting. Obtain medical attention immediately.

Call Drug and Toxicology Information Services (DATIS) on +263242307148/+263786100098 for more information.

NOTE TO PHYSICIAN:

Antidote: There is no specific antidote. Treat symptomatically.

EMPTY CONTAINER DISPOSAL

Invert the container over the spray tank or mixing tank. Allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water to a minimum of 25% of the container. Add the rinsate to the contents of the spray tank. Destroy the empty container by perforation and flattening. **DO NOT** use for any other purposes. If possible, offer the container for recycling and reuse.

CLEANING OF APPLICATION EQUIPMENT

After use, ensure traces of FIVESTAR 325 SC are removed. Use the following method:

- a) Drain and rinse tank, spray boom and hoses with clean water for at least 10 minutes (rinse at least 3 times).
- b) Fill tank with clean water and add to it 1 liter of household bleach (Sodium hydrochlorite, 5%), or 1.5-liter household bleach 3.5% per 200 liters of water. Rinse hoses and spray boom and leave in tank for 15 minutes whilst agitating. Drain through nozzle outlets.
- c) Repeat step (b) and after that, rinse thoroughly with water. Dispose of the wash water safely in a collection sump where it will not contaminate crops, grazing, rivers, dams and boreholes.

WARRANTY:

The excellent results which this chemical normally yields are not however warranted or guaranteed as the effectiveness can be greatly influenced by factors outside the control of either the manufacturer or the distributor. No warranty whatsoever expressed or implied is therefore given concerning the effectiveness of this chemical. Any responsibility is specifically excluded for any damage, injury or loss of any kind of loss during or resulting from its handling, use or storage whether or not such has been in accordance with label directions.



FIVESTAR

325 SC

Reg. No.:
23-B-73-40-B-61-35

CONTENTS:
1L



Composition:.....	mass/volume
Azoxystrobin:.....	200 g/l
Difenoconazole:.....	125 g/l
Inert ingredients:.....	675 g/l

Chemical group:

Azoxystrobin:.....	strobilurin
Difenoconazole:.....	triazole

A broad spectrum suspension concentrate fungicide with systemic, translaminar and contact properties for the preventive control of different leaf diseases on crops as listed.

Manufactured by:
Shandong Weifang Rainbow Chemical Co., Ltd.
Binhai Economic Development Area, Weifang, Shandong, China
Tel: +86-531-88875222, 88875231

Distributed By:
Trowbar Enterprises (PVT) LTD. T/A MAGUIRES
196 Harare Drive, Borrowdale, Harare, Zimbabwe
+263 (0242) 852 131

Shelf Life: Two years from the date of manufacture if stored in a cool dry place in original unopened container.
Date of Manufacture:
Expiry Date:
Batch No.:



FUNGICIDE

DIRECTION FOR USE: USE AS DIRECTED.

MIXING INSTRUCTIONS

Half fill the spray container with clean water. Add the required amount of FIVESTAR 325 SC to the spray container and top up with water to the required volume while agitating continuously.

Use the prepared mixture immediately. **DO NOT** prepare more spray mixture than is needed for immediate application. **DO NOT** allow to stand overnight.

APPLICATION RATES:

The application must be repeated if it rains within two hours after application.

Use 100L water per hectare.

Crop & Pest	Dosage/ha	Application directions
Dry Beans Anthracnose, Rust	500 mL/ha 16L Knapsack sprayer: 80mL	Apply 2 consecutive sprays 14 days apart starting at flower initiation stage. DO NOT exceed two applications per season.
Groundnuts Early leaf spot, Web blight		Apply 2 consecutive sprays 14 - 21 days apart starting at first signs of disease, or at flower initiation (10 - 12 weeks after planting) if no disease is present. DO NOT exceed two applications per season.
Maize, Sweetcorn, and Grain Sorghum Grey leaf spot, Corn leaf blight, Rust	500 - 750mL/ha 16L Knapsack sprayer: 80-120mL	Start application before 3 % of the total leaf area shows disease symptoms OR as soon as symptoms appear on the basal 3 - 5 leaves. Apply at 21 to 28-day intervals. Use the shorter application interval (7 to 10 - days) if disease pressure is very high. DO NOT apply more than three sprays per season.
Potatoes, Tomatoes Early blight	500 mL/ha 16L Knapsack sprayer: 80mL	Apply in blocks of two consecutive sprays 7 - 10 days apart when the weather favours disease development. DO NOT exceed six applications per season.
Brassicas Alternaria, Anthracnose, Leaf spot, Powdery mildew	800-1000mL/ha 16L Knapsack sprayer: 128-160mL	Apply before disease occurs. Apply at 7-10 day intervals. DO NOT exceed four (4) applications per season.
Cucurbits Anthracnose, Leaf blight, Leafspot, Downy mildew, Powdery mildew	730 - 1000mL/ha 16L Knapsack sprayer: 120-160mL	Apply up to three (3) consecutive sprays 7 to 10 days apart starting at flower initiation. DO NOT apply PRODUCT with wetting agents. DO NOT make more than three (3) applications per season.
BULB VEGETABLES (Chives, garlic, leeks, onions) Botrytis leaf blight, Cercospora leaf spot, Powdery mildew, Purple blotch, Stemphylium leaf blight	500 - 1000 mL/ha 16L Knapsack sprayer: 80-160mL	Apply as part of a 7 to 10 days spray schedule. DO NOT apply more than one(1) application before following-up with products with a different mode of action. DO NOT exceed four (4) applications per season.

RE-ENTRY PERIOD

DO NOT enter the treated area for at least 24 hours after treatment.

COMPATIBILITY

Compatible with most of the commonly used pesticides and other crop protection products. However, all mixtures should be tested prior to field application.