

**BEFORE YOU OPEN THE CONTAINER,
READ AND UNDERSTAND THE ENTIRE LABEL.**

PRECAUTIONS:

- Handle with care; avoid splashing / contact; poisonous by swallowing, inhalation and skin contact. Do not smoke, eat or drink while using this product.
- Store in original container under LOCK AND KEY. KEEP OUT OF REACH OF CHILDREN. KEEP APART FROM FOOD AND FOODSTUFFS.
- MAY BE DANGEROUS TO ANIMALS, BIRDS, FISH AND BEES.
- DO NOT CONTAMINATE DRINKING POOLS, DAMS, RIVERS, WATERWAYS AND OTHER WATER SUPPLIES.

FOR SAFETY WHEN MIXING

- Wear eye protection (at least glasses). Undiluted Fusilade Forte® 150 EC can irritate your eyes. If you get it in your eyes, wash it out at once. See **FIRST AID**.
- Wear rubber gloves. Undiluted Fusilade Forte® 150 EC can irritate your skin. If you get it on your skin, wash it off at once. If you spill it on your clothes, change and wash them immediately.
- Triple rinse empty container and pour washings into spray tank.

FOR SAFETY DURING SPRAYING:

- As with all chemicals AVOID CONTACT WITH SPRAY as much as you can.
- Wear overalls and boots.

FOR SAFETY AFTER SPRAYING:

- Dispose of unwanted spray solution by spraying off on waste ground.
- Remove protective clothing and wash it before re-use.
- WASH YOURSELF with soap and water.
- Destroy empty container by crushing or cutting then place in secure disposal site - DO NOT USE for any other purpose.
- Clean applicator thoroughly. Dispose of wash water in a safe manner.

SYMPTOMS OF HUMAN POISONING: No known symptoms of poisoning.

FIRST AID: Eye splashes: Hold eyelids apart and pour in a gentle stream of water for 10-15 minutes. Go to a doctor.

If Fusilade Forte® 150 EC is SWALLOWED do not make the person vomit. Take person, and this label, to a DOCTOR AT ONCE.

NOTE TO PHYSICIAN: Perform gastric lavage, taking care to prevent aspiration of gastric contents. Consider administration of activated charcoal and a laxative. Treat symptomatically.

© Registered Trademark of a Syngenta Group Company
© 2005 Syngenta Crop Protection AG

L165274 ZIMB/05Q PPE 4012228

 **Fusilade Forte®**
150EC

syngenta®

Reg. No.:
05-C-107-4

Net Content:
1 Litre

**TO CAUSE A HAZARD IN
THE USE, STORAGE OR
DISPOSAL OF THIS SUB-
STANCE IS AN OFFENCE.**


Shelf Life: A minimum of 2 years
if stored in a cool, dry place in
original unopened container.

Registered by: 119 MM
Syngenta Agro Services
P.O. Box 1088, Harare
Suite Za, Julia's Parade
Sam Levy, Borrowdale
Tel.: 0772 402 477

Emergency Call Number:
Syngenta Alarm Centre
in Huddersfield, UK
Tel.: +44 1484 538 444
Fax: +44 1484 554 093

Manufactured by:
Syngenta Crop Protection AG
Basle Switzerland

Product names marked ® or TM, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company



HARMFUL IF SWALLOWED

COMPOSITION (mass/vol)
fluzifop-p-butyl 150 g /litre
inert ingredients to 1 litre
Chemical group:
Aryloxyphenoxy propionate

A post emergent selective translocated herbicide for the control of most annual and perennial grass weeds as well as the suppression of upright starbur in tobacco, cotton, groundnuts, beans and soya-beans. A sugar cane ripener.



CAUTION



AREA OCCUPIED BY LEAFLET

MAXIMUM TYPE AREA
58 MM X 119 MM

S-PAC

DESCRIPTION: LBL LFT S FANFOLD
8 PG RH 270X122 MM

SCALE:
1:1

ISSUE NO:
SYN002

ISSUE DATE:
03.08.2009

234232

APPLICATION	RATE	COMMENT
Early season	275 ml / ha	The time of application is related to the date of harvest (i.e. 6-10 weeks prior to the set harvesting date) however, application also depends on cane maturity.

PAGE 6

WARNING:

Spray drift or direct contact with young cane can result in crop damage.

66mm x 117mm

WARRANTY:

All products supplied by Syngenta Agrochemicals are to be used only in accordance with the recommendations and instructions given on the labels provided with this pack. Further, although it is believed that all products supplied by us are of high quality and are suitable, for the purposes indicated on the label, since we cannot exercise control over the every day use of products, we do not give any warranty whatsoever as to their quality and fitness for purposes and we will not accept any responsibility whatsoever for damage arising from the storage, handling, application or use of our products.

6

Emergency Call Number:
 Syngenta Alarm Centre in Huddersfield, UK
 Tel.: +44 1484 538 444
 Fax: +44 1484 554 093

Manufactured by:
 Syngenta Crop Protection AG
 Basle, Switzerland

© 2005 Syngenta Crop Protection AG

® Registered Trademark of a Syngenta Group Company

PAGE 7

BACK PAGE

MAXIMUM TYPE AREA
 54.5mm x 117mm

8

L165274 ZIMB/05Q PPE 4012228

Fusilade Forte
 150 EC

750ml

BEFORE YOU OPEN THE CONTAINER, READ AND UNDERSTAND THE ENTIRE LABEL.

approx. measure

Reg. No.: 05-C-107-4

COMPOSITION (mass/vol)
 fluazifop-p-butyl 150 g / litre
 inert ingredients to 1 litre
 Chemical group: Aryloxyphenoxy propionate

500ml

A post emergent selective translocated herbicide for the control of most annual and perennial grass weeds as well as the suppression of upright starbur in tobacco, cotton, groundnuts, beans and soyabeans. A sugar cane ripener.

DIRECTIONS FOR USE

1. Fusilade Forte® 150 EC AS A HERBICIDE:
 Fusilade Forte® 150 EC may be used to control most annual and perennial grass weeds as well as the suppression of Upright Starbur in Burley and Virginia tobacco in the lands, cotton, groundnuts, beans and soyabeans.

250ml

S-PAC

DESCRIPTION: LBL LFT S FANFOLD
 8 BC RH 270X122 MM

SCALE:
 1:1

ISSUE NO:
 SYN002

ISSUE DATE:
 03.08.2009

234232

COUNTENANCED BY TOBACCO RESEARCH BOARD OF ZIMBABWE.

CERTIFICATE NO: 12-15-C-12

PRECAUTIONS:

- Handle with care; avoid splashing / contact; poisonous by swallowing, inhalation and skin contact. Do not smoke, eat or drink while using this product.
- Store in original container under LOCK AND KEY. KEEP OUT OF REACH OF CHILDREN. KEEP APART FROM FOOD AND FOOD-STUFFS.
- MAY BE DANGEROUS TO ANIMALS, BIRDS, FISH AND BEES.
- DO NOT CONTAMINATE DRINKING POOLS, DAMS, RIVERS, WATERWAYS AND OTHER WATER SUPPLIES.

FOR SAFETY WHEN MIXING

- Wear eye protection (at least glasses). Undiluted Fusilade Forte® 150 EC can irritate your eyes. If you get it in your eyes, wash it out at once. **See FIRST AID.**
- Wear rubber gloves. Undiluted Fusilade Forte® 150 EC can irritate your skin. If you get it on your skin, wash it off at once. If you spill it on your clothes, change and wash them immediately.
- Triple rinse empty container and pour washings into spray tank.

FOR SAFETY DURING SPRAYING:

- As with all chemicals AVOID CONTACT WITH SPRAY as much as you can.
- Wear overalls and boots.

FOR SAFETY AFTER SPRAYING:

- Dispose of unwanted spray solution by spraying off on waste ground.
- Remove protective clothing and wash it before re-use.
- WASH YOURSELF with soap and water.
- Destroy empty container by crushing or cutting then place in secure disposal site - DO NOT USE for any other purpose.
- Clean applicator thoroughly. Dispose of wash water in a safe manner.

SYMPTOMS OF HUMAN POISONING: No known symptoms of poisoning.

FIRST AID: Eye splashes: Hold eyelids apart and pour in a gentle stream of water for 10-15 minutes. Go to a doctor.

If Fusilade Forte® 150 EC is SWALLOWED do not make the person vomit. Take person, and this label, to a DOCTOR AT ONCE.

NOTE TO PHYSICIAN: Perform gastric lavage, taking care to prevent aspiration of gastric contents. Consider administration of activated charcoal and a laxative. Treat symptomatically.

RATES

CROP STAGE	WEEDS/WEED HEIGHT	RATE L / HA
Tobacco Seedbeds	Annual grasses	0.58
Field (tobacco, cotton, groundnuts, beans and soyabeans)	Annual grasses: < 120 mm 120-250 mm > 250 mm	0.830 1.25 1.7-2.1
	Mature perennial grasses: Undisturbed rhizomes Fragmented rhizomes	2.5-6.7 1.25-2.9
	Shamva and sorghum volunteers (up to 200 mm height)	0.4
	Wheat and barley volunteers (up to 5 leaf stage)	1.25

APPLICATION: A relatively short contact period of 1.5-2 hours without rain is required for satisfactory results.

Tobacco: Fusilade Forte® 150 EC can be directed onto grass weeds or applied as a broadcast spray anytime after transplanting, until within 3 weeks of harvesting.

Fusilade Forte® 150 EC contains a wetter. No additional wetter is required. Fusilade Forte® 150 EC may be applied through any sprayer fitted with normal fan, cone or flood jet nozzles. For ground application 100-400 litres spray / ha should be

applied. For aerial application 20-30 litres spray / ha should be applied. Clean water should always be used.

HARVEST INTERVAL (Providing rate has not exceeded 3.0 litres/ha):
Cotton, Groundnuts, Phaseolus beans and Soyabeans 8 weeks
Tobacco 3 weeks.

WARNINGS: Do not plant any of the following crops: maize, sorghum, wheat, barley, or grasses within 2 months of a Fusilade Forte® 150 EC application. With Fusilade Forte® 150 EC, weeds should be actively growing and not stressed, at the time of application.

COMPATIBILITY: It is not recommended to tank mix Fusilade Forte® 150 EC with other materials. These should be sprayed in a sequence of at least five days apart. The order in which Fusilade Forte® 150 EC and other products are applied is not important.

2. Fusilade Forte® 150 EC AS A SUGAR CANE RIPENER:

APPLICATION DIRECTIONS AND RATES: Fusilade Forte® 150 EC should be applied to EARLY season sugar cane harvests (April-June harvests).