


BASF

Kumulus® DF

Registration No.:
96-B-3-14

25 kg net



Harmful if swallowed

Composition	mass/mass
sulphur	80 %
inert ingredients	20 %

An Efficient Fungicide for the Control of Powdery Mildew and other Fungi. Also controls certain Mite Species.

Manufactured by: BASF Aktiengesellschaft

Registered by: Pivotal Agro-Service (Pvt) Ltd.
P.O. Box UA 414, Union Ave, Harare, Telephone 31 01 62-4,
Telefax 33 11 68, e-mail: info@basf.co.zw

® = Registered trademark of BASF

84188 7/ZW 0082

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE**PRECAUTIONS**

1. Handle with care; avoid splashing/contact with skin.
2. KEEP OUT OF REACH OF CHILDREN.
3. KEEP APART FROM FOOD AND FOODSTUFFS.
4. Store in original container and KEEP UNDER "LOCK AND KEY".
5. Do not eat, drink or smoke, while applying this fungicide.
6. Only prepare as much spray solution as necessary for immediate use.
7. Empty container thoroughly into the spray tank and dispose of safely.
8. Destroy empty container and DO NOT use for any other purpose.
9. DO NOT CONTAMINATE PONDS, WATERWAYS, DRINKING POOLS OR DITCHES WITH THE CHEMICAL.

SYMPTOMS OF POISONING

Ingestion of large amounts may lead to the following:

Severe mucous membrane and gastrointestinal irritation with nausea, vomiting and diarrhoea. Central nervous system depression with respiratory depression, headache, dizziness, paleness, rapid, feeble pulse, cyanosis, low blood pressure, loss of consciousness, tremors, convulsions, coma, death from respiratory paralysis.

FIRST AID

1. If swallowed, call medical help and show this label. Milk can be given while waiting for medical help.
2. Eye contamination: wash with copious amounts of water.
3. Skin contamination: wash with soap and water.
4. If inhaled, keep patient calm, remove to fresh air.

NOTE TO PHYSICIAN

If swallowed, give milk, induce emesis (unless tissue erosion evident) or perform careful gastric lavage (avoid perforation). Administer gastric antacids and demulcents. Administer artificial respiration and oxygen. If excessive vomiting or diarrhoea occurs, maintain fluid and electrolyte balance.

WARNINGS

1. DO NOT apply to cucumbers or sweet melons.
2. DO NOT spray Kumulus DF in very hot weather or should spraying be essential at such times, endeavour to spray during the cooler parts of the day.
3. DO NOT apply Kumulus DF four weeks before or after application of an oil or oil-based spray.

WARRANTY

Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use are beyond our control and we are unable to foresee all circumstances arising therefrom, we do not accept any liability for damage which may result from the storage and use of the product. We are liable for the consistent quality of the product, but the risk of its storage and use is not borne by us.

DIRECTIONS FOR USE

CROP	DISEASE/PEST	RATE/100 L WATER	APPLICATION
APPLES	Powdery mildew	300 g	Apply a good cover spray during the critical October/November bud movement period at weekly intervals. Six applications may be required. DO NOT apply to sulphur-shy varieties such as Golden Delicious as russetting may occur.
CITRUS	Concentric ring blotch	150-200 g	Spray as a light cover spray ensuring complete and uniform wetting of all leaves and fruits. Make the first application as soon after petal fall as possible and repeat 2-3 months later. Sprays should in any case be timed before disease symptoms appear.
	Grey mites and Brevipalpus mites	150-200 g	Spray as a light cover spray ensuring complete and uniform wetting of all leaves and fruits. Spray only after scouting indicates the presence of mites on the trees.
CUCURBITS (NOT Cucumbers or sweet melons)	Powdery mildew	300 g	For use on pumpkins and squash. Start spraying at first sign of disease and continue, where necessary at 7-10 day intervals.
GROUND-NUTS	Cercospora Leaf spot	200 g	Start spraying at the first sign of disease and repeat after each heavy rain or at 14 day intervals.
PEACHES NECTARINES APRICOTS	Powdery mildew	200-300 g	Apply a good cover spray starting at 75 % petal-fall stage and continue with 2-3 further precautionary sprays at 14-21 day intervals. Additional sprays may be necessary in orchards where severe mildew attacks can be expected.
PEAS	Powdery mildew	150-200 g	Start spraying at the first sign of disease and continue at 14 day intervals. With overhead irrigated crops, apply as soon as practical after irrigation.
TOMATOES	Powdery mildew	200 g	Start spraying at the first sign of disease and repeat at 7 day intervals. This spraying programme will also control russet mite.
VINES	Powdery mildew	200 g	Kumulus DF in conjunction with Polyram DF will provide a spray programme for season-long control of disease. Kumulus DF will control powdery mildew (Uncinula necator). Spray to run off. First spray when the shoots are 10-20 cm long, and the second when shoots are 30-35 cm long. These first two applications are important and are applied irrespective of weather conditions. Continue spraying at 14 day intervals or after heavy rain. Stop spraying 4-6 weeks before harvest as the spray may tarnish the berries.

COMPATIBILITY

Kumulus DF is compatible with most commonly used insecticides and fungicides and, in particular, is fully compatible with Polyram DF.


DO NOT mix with mineral oils, oil based sprays, such as DNOC Peach and Winter oils.

BASF

Kumulus® DF

Registration No.:
96-B-3-14

25 kg net



Harmful if swallowed

Composition	mass/mass
sulphur	80 %
inert ingredients	20 %

An Efficient Fungicide for the Control of Powdery Mildew and other Fungi. Also controls certain Mite Species.

Manufactured by: BASF Aktiengesellschaft

Registered by: Pivotal Agro-Service (Pvt) Ltd.
P.O. Box UA 414, Union Ave, Harare, Telephone 31 01 62-4,
Telefax 33 11 68, e-mail: info@basf.co.zw

® = Registered trademark of BASF

84188 7/ZW 0082



Batch No.

Date of manufacture:

Expiry date: 2 years after date of manufacture



4 014348 841887