

L1085335 ZIMB/01A PPE 4153945



Maxim[®] XL
035FS

syngenta.

Before using this product read and understand the entire label.

**TO CAUSE A HAZARD IN THE USE,
STORAGE OR DISPOSAL OF THIS SUBSTANCE
IS AN OFFENCE.**

Shelf-life: Two years from date of manufacture if stored in original container under constant cool conditions.

Date of manufacture: / Batch No:

Please refer to Inkjet print.

Emergency Call Number:

Swiss Toxicological Information Centre
(24 h) +41 44 251 51 51

Manufacturer:

Syngenta Crop Protection AG, Basel, Switzerland

® Registered Trademark of a Syngenta Group Company

Registered by: Syngenta Agro AG

32 Sandringham Drive, Alexandra Park, Harare.
Tel.: 08677005432/08677005434/0772402477

5 Litres

Product names marked ® or TM, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company



PRECAUTIONS

1. Handle with care; avoid splashing / contact; poisonous by swallowing, inhalation and contact with skin.
2. DANGEROUS TO FISH.
3. KEEP OUT OF REACH OF CHILDREN.
4. KEEP APART FROM FOOD, FOODSTUFFS, seeds and fertilizers.
5. Store in original container in a cool, dry place and KEEP UNDER LOCK AND KEY.
6. Use only on the crops for which this product is registered.
7. Decontamination of Mixing Tank – clean the tank thoroughly after use and ensure that all traces of MAXIM® XL 035 FS are removed. Make use of the following method:
(a) Drain tank and rinse tank and hoses with clean water for at least 10 minutes. (b) Fill tank with clean water and add to it 1,0 litre household bleach (5%) or 1,5 litres household bleach (3,5%) per 200 litres of water. Rinse hoses and leave in the tank for 15 minutes whilst agitating. (c) Repeat step (b) and thereafter, rinse thoroughly with clean water and dispose of the wash water at a site designated for the disposal of pesticides.
8. Empty container disposal – invert the empty container over the mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container 3 times with a volume of water equal to at least 10% of that of the container. And the rinsate to the contents of the mixing tank. Destroy the empty container by perforation and flattening. Place it in a secure disposal area and offer it for recycling. DO NOT use it for any other purpose.

SYMPTOMS OF POISONING

No case of human poisoning is on record.

FIRST AID

If poisoning is suspected, stop work immediately and seek medical advice. In case of contact with skin and eyes, wash immediately with plenty of water. Rinse splashes from eyes with abundant fresh and clean water for at least 10 minutes. If inhaled move to clean air. If swallowed, give medical charcoal in plenty of water.

NOTE TO PHYSICIAN

No specific antidote is known. Apply symptomatic therapy.

WARRANTY

Before applying the product read the entire label.

WARNING

Maxim® XL 035 FS contains Metalaxyl-M. Experience has shown that strains of fungus resistant to Metalaxyl-M may develop, resulting in a reduction in the efficacy of the product and thus yield losses. Since the occurrence of resistance cannot be foreseen, the manufacturer accepts no responsibility for any loss of, or damage to, crop resulting from failure of Maxim® XL 035 FS to control the resistant strains.

USER'S RISK

The user bears the risk for damage resulting from factors beyond the manufacturer's control.

Resistant strains of fungal diseases may develop or may exist against which fungicides may not be effective resulting in crop loss. Since the occurrence of such strains cannot be forecast, neither the manufacturer nor its distributors can accept responsibility for any loss or damage caused by failure of fungicides to control resistant strains.

All recommendations for use of the product are based on the current state of the manufacturer's knowledge. Since the manufacturer cannot control the application, use or storage of the product, the manufacturer cannot accept responsibility therefore.

SPECTRUM OF ACTIVITY

Fludioxonil controls economically important diseases caused by fungi in the classes of Ascomycetes, Basidiomycetes and Fungi imperfecti.

Metalaxyl-M controls economically important diseases caused by fungi in the class of Oomycetes.

CROP TOLERANCE

When used as recommended, Maxim® XL 035 FS is well tolerated by the seed and the seedling. In case of doubt, check crop tolerance first.

DIRECTIONS FOR USE

Crop	Target Pathogen	Product rate: ml/100 kg seed
Maize	<i>Fusarium, Pythium</i>	100
Sorghum	<i>Fusarium, Pythium</i>	100
Cotton	<i>Rhizoctonia, Pythium</i>	100
Peas	<i>Ascochyta, Pythium</i>	100
Beans	<i>Fusarium, Pythium</i>	100
Soyabeans	<i>Fusarium, Pythium</i>	100
Sunflowers	<i>Downy mildew, Fusarium</i>	100

APPLICATION TECHNIQUE

Maxim® XL 035 FS can be applied on any continuous flow seed treater if the formulation is sprayed onto the seed through nozzles or spinning disks in a liquid mist chamber. Rotating drums or fluidized bed systems for film coating procedures are also recommended. It is essential that any continuous flow seed treater should be calibrated with Maxim® XL 035 FS, as flow characteristics may not be the same as for other seed treatment products.

When used diluted, the following procedure should be followed for preparing the slurry:

- Fill the mixing tank with water.
- Shake the product before use.
- Add the required quantity of Maxim® XL 035 FS while stirring continuously.
- Maintain agitation of the slurry during the whole time of use.
- The slurry has to be used within 24 hours after preparation.

TREATMENT PROCEDURE

Product + Water = Slurry (to be applied onto seeds)

100 ml + 700-1100 ml = 800-1200 ml/100 kg seed

Note: Clean equipment with water after use.

Calibrate the flow of seed in a seed drill or planter.

Treated seeds should be sown properly and completely covered by soil. Excess seed are not to be used as food for birds or other animals.

COMPATIBILITY

Maxim® XL 035 FS is compatible with film coating products and with FS/WS formulations of other seed treatment compounds with neutral reaction.

Registered by: Syngenta Agro AG

32 Sandringham Drive, Alexandra Park, Harare.

Tel.: 08677005432/08677005434/0772402477

Manufacturer:

Syngenta Crop Protection AG, Basel, Switzerland

Emergency Call Number:

Swiss Toxicological Information Centre

(24 h) +41 44 251 51 51



NO TEXT HERE
GLUE PAGE
97 x 134 mm