



300 SC

Zambia Reg. No. ZEMA/LSK/IP/011/2012
Reg. No. 12-B-60-19-B-72-3

HARMFUL IF SWALLOWED

COMPOSITION:

Tebuconazole	200 g/l
Trifloxystrobin.....	100 g/l
Inert ingredients.....	700 g/l

MASS/MASS

Chemical Group : Tebuconazole - Triazole
Trifloxystrobin - Strobilurin

NATIVO 300 SC IS A SYSTEMIC SUSPENSION CONCENTRATE FUNGICIDE WITH MESOSTEMIC PROPERTIES FOR THE CONTROL OF RUST IN COFFEE & SOYBEANS, EARLY BLIGHT IN POTATOES & TOMATOES, GREY LEAF SPOT IN MAIZE, LEAF RUST & POWDERY MILDEW IN WHEAT & BARLEY AND ALTERNARIA & FROG EYE IN TOBACCO

Packed For:
Bayer Zimbabwe (Pvt) Ltd
P.O. Box AY 78, Amby, Harare, Zimbabwe.
Tel. 487211/487245-6 Fax: 487242

IN CASE OF POISONING PLEASE PHONE: +27 (21) 931-6129

Manufactured by: Bayer CropScience AG

Nativo® is a registered trademark of Bayer CropScience AG, Germany

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.

PRECAUTIONS:

- Handle with care; avoid any contact, poisonous by swallowing or inhalation.
- Do not eat, drink or smoke while mixing, applying, handling or before washing hands and face.
- WEAR PROTECTIVE CLOTHING, i.e. overalls, rubber gloves, and rubber boots and a facemask while mixing, spraying and cleaning up.
- Do not inhale fumes or spray mist.
- Remove protective clothing on completion of spraying, wash thoroughly with soap and water, and wash contaminated clothing daily.
- Clean applicator after use, dispose of wash water where it will not contaminate crops, grazing, rivers and dams.
- KEEP APART FROM FOOD AND FEEDSTUFFS
- KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
- STORE IN A COOL, DRY PLACE IN ORIGINAL CONTAINER AND KEEP UNDER LOCK AND KEY.
- TOXIC TO FISH AND WILD LIFE. DO NOT CONTAMINATE DRINKING POOLS, DAMS, RIVERS AND WATERWAYS.
- SPILLAGE - Use protective clothing when dealing with spillages. Prevent entry into drains, water or soil. Soak up spilled product with absorbent material such as dry soil, sawdust or sand and place into a labeled closable container for safe disposal. Use damp cloth to clean the floor and contaminated parts. Place used cleaning materials into closable container for safe disposal. Bury container in a disposable pit away from water sources or arable land.
- CONTAINER DISPOSAL - Triple rinse container emptying washings into spray tank and spray onto crop. Perforate and flatten rinsed container to make it unusable or burn it and bury crushed container or ashes in a recommended disposal pit in a secure location away from water sources or arable land. Do not use empty container for any other purpose.
- R.T.:** 12 hours, but do not enter treated area before spray deposits on leaf surfaces have dried, unless wearing protective clothing.

SYMPTOMS OF POISONING:

No specific symptoms known.

FIRST AID:

- Remove patient from source of poisoning and remove contaminated clothing
- In case of skin contact, remove clothes and bathe with soap and water and take the patient to the doctor.
- If eyes are contaminated, flush well with clean water for at least 15 minutes, cover the affected eye and take the patient to the doctor.
- In case of poisoning by inhalation, remove the victim from the contamination zone, move outdoors and keep the patient lying sideways and monitor the breathing.
- Show container with label to attending doctor.

NOTE TO PHYSICIAN:

Treatment:

Treatment has to be symptomatic and supportive

General decontamination measures, if it has been ingested, gastric lavage practice.

Antidote: No specific antidote is known.

WARRANTY:

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label and the occurrence of resistance of the pathogen against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

DIRECTIONS FOR USE: Use only as directed.

MIXING INSTRUCTIONS:

Add the required amount of **Nativo 300 SC** into spray tank half-filled with clean water with the agitation system in operation and then top up with water to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.

CROP	DISEASE	DOSAGE	REMARKS
COFFEE	Rust <i>Hemileia vastatrix</i>	750 – 1000 m ^l /ha Apply at 750-1000 m ^l Nativo 300 SC in 1000-2000 ℓ water/ha	Spray at first sign of infection. Repeat as necessary after 3-4 weeks. P.H.I. Pre-harvest interval: 30 days.
POTATOES & TOMATOES	Early blight <i>Alternaria solani</i>	Ground application: 800 m ^l /ha in 500 ℓ water/ha.	Apply a series of maximum 3 sprays in a preventive programme at 7 day intervals starting about 4 weeks after full emergence (tuber initiation/flowering). NB. Nativo 300 SC should be applied in a tank mixture with a registered late blight fungicide such as Infinito or Melody duo when conditions are favourable for the development of late blight. P.H.I. Pre-harvest interval: 7 days.
SOYBEANS	Rust <i>Phakopsora pachyrhizi</i>	Ground application: 500 m ^l /ha. Apply 500 m ^l Nativo 300 SC in 200-300 ℓ water/ha Aerial application: 500 m ^l /ha applied in 30-40 ℓ water/ha	Apply as a protective programme. 1st application: Apply at early flowering (approx. 50 days after planting) 2nd application: Apply at 20 day interval P.H.I. Pre-harvest interval: 30 days.
MAIZE	Grey leaf spot <i>Cercospora zeae-maydis</i>	1000 m ^l /ha Apply 1000 m ^l Nativo 300 SC in 200 – 300 ℓ water/ ha	Apply when disease symptoms are first observed on the basal 4-5 leaves and before the infection level on the lowest leaves reaches 5 % and before lesions appear on the 3 leaves below the cob (approximately 55-75 days after planting). Follow-up with a second application 21-25 days later.
WHEAT & BARLEY	Leaf rust <i>Puccinia</i> sp. Powdery mildew <i>Erysiphe graminis</i>	Ground application: 500 m ^l /ha applied in 250 – 300 ℓ water/ha. Aerial application: 500 m ^l /ha applied in 30 – 40 ℓ water/ha.	Commence spraying at the very first signs of the disease / infection. Apply the 2nd application 3 - 4 weeks later.
TOBACCO Lands Countenanced by Tobacco Research Board (TRB) Certificate No.	Alternaria (<i>Alternaria alternata</i>) and Frogeye (<i>Cercospora nicotianae</i>)	Ground application: 600 m ^l /ha applied in 200 water/ha.	Start spraying at 8 weeks after planting continuing at 14 day intervals up to 14 weeks after planting, applying a maximum of 4 sprays in any given season. Spray method: High volume - 200 ℓ water/ha

P.H.I. Pre-harvest interval: time between application and harvest.

R.T. Re-entry time: time after application and re-entry into treated area.

