

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE

PRECAUTIONS:

1. Handle with care and avoid any contact. Poisonous by swallowing or inhalation.
2. **Oberon** may cause sensitization by skin contact.
3. Do not eat, drink or smoke while mixing, applying, handling or before washing hands and face.
4. WEAR PROTECTIVE CLOTHING, i.e. overalls, rubber gloves, and rubber boots and a facemask while mixing, spraying and cleaning up.
5. Do not inhale mist or spray against the wind.
6. Remove protective clothing on completion of spraying, wash thoroughly with soap and water, and wash contaminated clothing daily.
7. Clean applicator after use, dispose of wash water where it will not contaminate crops, grazing, rivers and dams.
8. KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
9. STORE IN A COOL DRY PLACE, IN THE ORIGINAL CONTAINER AND KEEP UNDER LOCK AND KEY.
10. KEEP APART FROM FOOD AND FOODSTUFFS.
11. TOXIC TO FISH AND WILD LIFE. DO NOT CONTAMINATE DRINKING POOLS, DAMS, RIVERS AND WATERWAYS.
12. SPILLAGE - Use protective clothing when dealing with spillages. Prevent entry into drains, water or soil. Soak up spilled product with absorbent material such as dry soil, sawdust or sand and place into a labelled closable container for safe disposal. Use damp cloth to clean the floor and contaminated parts. Place used cleaning materials into closable container for safe disposal. Bury container in a disposable pit away from water sources or arable land.
13. CONTAINER DISPOSAL - Triple rinse container emptying washings into spray tank and spray onto crop. Perforate and flatten rinsed container to make it unusable and bury it in a disposal pit in a secure location away from water sources or arable land. Do not use empty container for any other purpose.
14. **RE-ENTRY:** Do not enter treated area before spray deposits on leaf surfaces have dried, unless wearing protective clothing.

SYMPTOMS OF POISONING: No specific symptoms known.

FIRST AID:

1. Remove patient from source of poisoning and remove contaminated clothing.
2. If skin is contaminated, wash affected areas of skin thoroughly with soap and water.
3. If eyes are contaminated, flush well with clean water for at least 15 minutes.
4. If swallowed, DO NOT induce vomiting, seek medical advice immediately.
5. Show container with label to attending doctor.

NOTE TO PHYSICIAN: Treatment: Treat symptomatically.

If ingested - removal of ingested compound by gastric lavage. No specific antidote is known

WARRANTY: Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest to the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.



o.b.e.r.o.n

Reg. No. 09-D-183-1

CONTENTS: 1L

CAUTION

HARMFUL IF SWALLOWED

COMPOSITION:

spiromesifen..... 240 g/l
inert ingredients..... 760 g/l

MASS/VOLUME

Chemical Group : ketoenol

A CONTACT ACARICIDE AND INSECTICIDE FOR THE CONTROL OF MITES AND WHITEFLIES ON ROSES, TOMATOES, POTATOES AND HYPERICUM



DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Oberon is a foliar contact acaricide and insecticide for the control of Red spider mites and Whiteflies on Roses, Tomatoes, Potatoes and Hypericum. **Oberon** is active against all developmental stages of mites and whiteflies, but the eggs, larvae, nymphs and pupae stages are more susceptible than adults.

Resistance management guidelines: **Oberon** may be applied up to 3 times in total per cropping cycle in annual crops. For mite control, a maximum of 2 applications, for whitefly, a maximum of 3 applications. **Oberon** applications should be alternated with registered products of other chemical groups with different mode of action.

Application: **Oberon** should be applied as a full cover spray on all foliage parts, ensuring a thorough and even coverage of upper and lower leaf surfaces. Apply a minimum overall spray volume of 1000 l of water/ha

Crop tolerance: **Oberon** is well tolerated by crops listed on the label, when used as recommended. **Note:** As the different species and varieties of Roses may differ in their sensitivity to chemical spray, users are advised always to check for crop compatibility by first treating a few plants before large-scale application is undertaken. Applications early in the morning or late in the afternoon when temperatures are lower and the drying time is longer may not show symptoms of selectivity but applications done at higher temperatures and low relative humidity may show symptoms of selectivity.

Mixing instructions: Thoroughly shake the container and pour the required quantity of **Oberon** into the spray tank half-filled with water while agitating and then top up with water to the required level. Do not allow the **Oberon** mixture to stand any length of time before using.

CROP	PEST	APPLICATION RATE	APPLICATION RATE
Roses	Red Spider mites	60 ml / 100 l water and apply a minimum of 1000 l/ha	Apply Oberon when presence of mites is first noticed and repeat at 10-day intervals if necessary. Note: 1. Early applications of Oberon will prevent mite population build up. 2. For effective control of high mites infestation, choose product effective on adults followed by Oberon .
Hypericum, Potatoes and Tomatoes	Red Spider mites and Whiteflies	60 ml / 100 l water and apply a minimum of 1000 l/ha or 9 ml/ knapsack of 15 l capacity	Apply Oberon when presence of mites or whitefly is first noticed and repeat at 10-day intervals if necessary.

Pre-harvest interval (PHI): Potatoes & Tomatoes: 3 days

Compatibility: **Oberon** should not be tank mixed with systemic and translaminar products. Do not mix **Oberon** with EC formulations or adjuvant. Do not apply EC formulations before or soon after **Oberon** application. **Oberon** is most efficacious when used alone at recommended rates and timing.

Packed For: Bayer Zimbabwe (Pvt) Ltd P.O. Box AY 78, Amby, Harare, Zimbabwe.
Tel. 487211/487245/7 Fax. 487242 Emergency Tel. +27(21) 931-6129
Manufactured by Bayer CropScience AG.

Oberon® is a registered trademark of Bayer CropScience AG, Germany

Batch No.: See on pack
Date of Manufacture: See on pack
Expiry date:



Zim/07/09/Oberon 240 SC/Full text label Code: 79661206



CAUTION

