

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE
PRECAUTIONS

- 1.Handle with care. Avoid splashing and contact with eyes, nose and mouth.
- 2.Poisonous by swallowing, inhalation and contact with skin.
- 3.**Wear full personal protective equipment (PPE)** i.e. rubber gloves, rubber boots, overalls, hood, goggles and respirator when handle with this product.
- 4.**KEEP APART FROM FOOD AND FOOD STUFFS.**
- 5.**DO NOT** eat, drink or smoke while preparing and applying this fungicide.
- 6.KEEP OUT OF REACH OF CHILDREN AND UNIFORMED PERSONS.
- 7.Dangerous to livestock and domestic pets.
- 8.Store in the original container and keep under lock and key.
- 9.Do not contaminate drinking pools, screams, water bodies with the chemical or empty containers.
- 10.VERY TOXIC TO AQUATIC ORGANISMS AND MAY CAUSE LONG-TERM ADVERSE EFFECTS IN THE QUATIC ENVIRONMENT.
- 11.Prevent drift onto other crops and areas not under treatment.

SYMPTOMS OF POISONING:

No specific symptoms of poisoning.

FIRST AID:

Inhalation: Move victim to fresh air. If not breathing properly, give artificial respiration. If breathing is difficult, give oxygen. Consult medical personnel.

Skin contact: Wash thoroughly with soap and running water. Remove contaminated clothing and shoes and launder.

Eye contact: Flush immediately with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

Ingestion: If swallowed, wash the mouth with water provided person is conscious. Induce vomiting. Obtain medical attention immediately.

Call Drug and Toxicology Information Services (DATIS) on +263242307148/+263786100098

NOTE TO PHYSICIAN:

Antidote: Administration of a gastric lavage with activated charcoal is recommended. Atropine sulphate should only be administered in case of obvious cholinesterase inhibition. Treatment should otherwise be supportive and symptomatic after appropriate decontamination. Treat symptomatically.

EMPTY CONTAINER DISPOSAL

Invert the container over the spray tank or mixing tank. Allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water to a minimum of 10% of the container. Add the rinsate to the contents of the spray tank. Destroy the empty container by perforation and flattening. Bury away in a designated area or away from water sources. DO NOT use for any other purposes. If possible, offer the container for recycling and reuse.

CLEANING OF APPLICATION EQUIPMENT

After use, ensure traces of Panol 722 SL are removed. Use the following method:


- a) Drain and rinse tank, spray boom and hoses with clean water for at least 10 minutes (rinse at least 3 times).
- b) Fill tank with clean water and add to it 1 liter of household bleach (Sodium hydrochlorite,5%) per 200 liters of water. Rinse hoses and spray boom and leave in tank for 15minutes whilst agita ng. Drain through nozzle outlets.
- c) Repeat step [b] and after that, rinse thoroughly with water. Dispose of the wash water safely in a collection sump where it will not contaminate crops, grazing, rivers, dams and boreholes.

WARRANTY:

The excellent results which this chemical normally yields are not however warranted or guaranteed as the effectiveness can be greatly influenced by factors outside the control of either the manufacturer or the distributor. No warranty whatsoever expressed or implied is therefore given concerning the effectiveness of this chemical. Any responsibility is specifically excluded for any damage, injury or loss of any kind of loss during or resulting from its handling, use or storage wether or not such has been in accordance with label directions.

Panol 722 SL

Reg. No.:
23-B-47-12
TRB No.:
25-26-B-120



HARMFUL IF SWALLOWED

CONTENTS:
1L

Composition:.....	mass/volume
Propamocarb Hydrochloride:.....	722 g/l
Inert ingredients:.....	278 g/l


Chemical group: Carbamate

A systemic fungicide for the prevention of damping-off diseases in agricultural and horticultural crops.

Manufactured by:
Shandong Weifang Rainbow Chemical Co., Ltd.
Binhai Economic Development Area, Weifang, Shandong, China
Tel: +86-531-88875222, 88875231

Distributed By:
Trowbar Enterprises (PVT) LTD. T/A MAGUIRES
196 Harare Drive, Borrowdale, Harare, Zimbabwe
+263 (0242) 852 131

Date of Manufacture:
Expiry Date:
Batch No.:



DIRECTION FOR USE:

USE AS DIRECTED.

MIXING INSTRUCTIONS

Half fill the spray container with clean water. Add the required amount of Panol 722 SL to the spray container and top up with water to the required volume while agitating continuously.

Use the prepared mixture immediately. Do not prepare more spray mixture than is needed for immediate application. Do not allow to stand overnight.

APPLICATION RATES:

Crop	Disease	Dosage/ha	Application directions
Tobacco seedbeds	Pythium root rot	4 ml/L water/ tray as a drench	Panol 722 SL should be applied before or at the early stage of the crop's disease occurrence.
Ornamentals and Flowers	Damping-off	75ml/ha in 300L water	Drench: Apply to seedbeds immediately after sowing at 30L of solution per hectare. Repeat at 2-3 weeks intervals
Avocado Seedlings in sleeves /Pots and Transplants	Root rot	Pots: 0.2-0.4ml/pot	20 ml dilute in 10L water. Drench Pot/Sleeves: with 100-200 ml of solution. Direct to stem base. Repeat at 3-4 weeks intervals with last treatment applied a week or less before transplanting.
		Lands: 800ml/ha in 400L water	Drench Post transplantat 400L of solution per hectare of soil surface in the planting basin. Repeat every 4 weeks.
Potatoes	Late blight (Phytophthora infestans)	1.4L/ha in 200-400L water	Apply Panol 722 SL on a 7-10 day spray interval, depending on disease pressure. The first spray should be applied as soon as there is a risk of blight infection. Maximum six treatments per year. Preventative treatment. Do not apply as a curative treatment when Late blight is present in the crop.

PHI

Ornamentals and Flowers 14days
Avocado 14-21days
Potatoes 14days

RE-ENTRY PERIOD

DO NOT enter the treated area for at least 24 hours after treatment.

COMPATIBILITY

Compatible with most of the commonly used pesticides and other crop protection products. However all mixtures should be tested prior to field application.