

SYMPTOMS OF POISONING:

Product may cause skin and eye irritation.

FIRST AID:

If on skin: Wash with soap and water. Get medical attention if irritation persists.

If in eyes: Flush eyes with clean water. Call a Physician if irritation persists.

If swallowed: Wash mouth with water if person is conscious. Call a Physician.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration and get medical attention.

NOTE TO PHYSICIAN:

There is no known antidote for exposure to this material. Treatment of overexposure should be directed at the control of symptoms and clinical condition.

WARRANTY:

This product and the recommendations concerning its use should give excellent results. Since the performance and effectiveness of crop protection chemicals can be greatly influenced by factors outside the manufacturer's control, the manufacturer and seller give no guarantee or warranty, expressed or implied concerning the use of this product and they assume no responsibility for damage or injury arising from the use thereof.

DIRECTIONS FOR USE

Apply as part of a **preventative spray program before the onset of diseases** in rotation with fungicides with a different mode of action. Apply in 300-1000 litres of water per hectare depending on area and type of crops.

COMPATIBILITY

Prestige 38 SC is compatible with most foliar fertilizers and insecticides such as BLANKET, FASTAC and VANTEX. For compatibility with other pesticides, consult respective labels or suppliers before tank-mixing with Prestige 38 SC.

WARNING:

Use product within 8 hours of mixing with water.

Do not apply more than 3 applications on one crop per season.

PRE-HARVEST INTERVALS: See recommendation table on leaflet, below.

RESTRICTED ENTRY INTERVAL: 24 Hours

RAINFASTNESS: 2 Hours

GENERAL INFORMATION

Prestige 38 SC is composed of active ingredients from two classes of fungicides, the strobilurin and anilid. **Prestige 38 SC** has multiple modes of action on most plant diseases and is effective against pathogens resistant to other fungicides. **Prestige 38 SC** is locally systemic, rapidly rainfast, inhibits spore germination, mycelial growth and sporulation of the fungus on the leaf surface. Optimum disease control is achieved when **Prestige 38 SC** is applied in a preventative program in rotation with a fungicide of a different mode of action like CUPERDEM.

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE**PRECAUTIONS:**

1. Handle with care and avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Harmful if inhaled or absorbed through the skin.
2. KEEP OUT OF REACH OF CHILDREN.
3. KEEP APART FROM FOOD AND FOODSTUFFS.
4. Remove contaminated clothing and wash before re-use.
5. Wash exposed parts of the body with soap and water after use and before eating, drinking or smoking.
6. **DO NOT CONTAMINATE PONDS, WATERWAYS AND DITCHES BY STORAGE OR DISPOSAL**
7. **DO NOT** eat, drink or smoke during application
8. **DECONTAMINATION OF SPRAYER**

Clean applicator/spraying equipment thoroughly after use and ensure that all traces of PRESTIGE 38 SC are removed. Use of the following method:

- a) Drain and rinse tank, spray boom and hoses with clean water for at least 10 minutes.
- b) Fill tank with clean water and add 1.0 litre household bleach (5%) or 1.5 litres of household bleach (3.5%) per 200 litres of water. Rinse hoses and spray boom and leave in tank for 15 minutes whilst agitating. Drain through nozzle outlets.
- c) Repeat step (b) and thereafter, rinse thoroughly with clean water and dispose of the wash water at a site designated for disposal of pesticides.

9. **EMPTY CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. Completely empty container into the application equipment and thereafter, thoroughly rinse the container 3 times with a volume of water equal to a volume of 10% of the container. Add the rinsings to the contents of the spray tank. Destroy empty containers by puncturing and burying. **DO NOT** use for any other purpose. If possible, offer for recycling.

10. **STORAGE:** Store in a cool, dry place in the original tightly closed container. **KEEP UNDER LOCK AND KEY.**

PRESTIGE 38 SC**REG. NO:**

19-B-88-7-B-90-3

Net Contents:

1 Litre

**HARMFUL IF SWALLOWED**

Pyraclostrobin.....12.8%

Boscalid.....25.2%

Chemical Group: Strobilurine + Carboxanilid

AVOCADO, BERRIES, BRASSICA, BULB/ROOT VEGETABLES, CARROTS, CITRUS, COTTON, CUCURBITS, FRUIT VEGETABLES, GRAPES, HERBS/FLOWERS, HOPS, LEAF GREENS, OIL SEEDS, PEAS/PULSES, POME FRUIT, POTATO, STONE FRUIT, TOMATO, TREE NUTS, TROPICAL FRUIT.

Registered by:

Pivotal Agro-Services (Pvt) Ltd

37 Kenmark Crescent

Bluff Hill Industrial Park

Harare, Zimbabwe

Telephone: +263 242 309825/6

Date of Manufacture: 06/06/2025

Batch Number: 2505014

PRESTIGE 38 SC APPLICATION RATES AND RECOMMENDATIONS:

CROP	DISEASE	RATE/HA	NO. OF SPRAYS	APPLICATION DIRECTIONS: Use only as directed.	PHI (Days)
Avocado	Anthracoese (<i>Colletotrichum gloeosporioides</i>) Scab (<i>Sphaeloma persea</i>)	800 ml	2	Commence application prior to the onset of disease development and repeat application 7-14 days later, as needed, or alternate with another registered fungicide with a different mode of action e.g. Cuperdem.	0
Berries: Blackberry Blueberry Gooseberry Raspberry Strawberry	Alternaria leaf spot (<i>Alternaria</i> spp.) Fruit rot (<i>Alternaria</i> spp.) Anthracoese (<i>Colletotrichum</i> spp.) Botrytis gray mold (<i>Botrytis cinerea</i>) Leaf spot and blotch (<i>Septoria</i> spp.) Monilinia blight (<i>Monilinia</i> spp.) Mummy berry (<i>Monilinia</i> spp.) Twig blight, and fruit rot (<i>Phomopsis</i> spp.) Downy mildew (<i>Plasmopara viticola</i>) Powdery mildew (<i>Microsphaera</i> spp., <i>Oidium</i> spp. and <i>Sphaerotheca</i> spp.)	300 – 500 ml	4	Commence applications of Prestige 38 SC prior to onset of disease and continue on a 7-day to 14-day interval. Use the shorter interval and/or the higher rate when disease pressure is high. To manage the potential for development of resistance, DO NOT make more than two sequential applications of Prestige 38 SC. Alternate with registered fungicide with a different mode of action. DO NOT apply Prestige 38 SC to blueberries as a tank mix with other pesticides, adjuvants, liquid fertilizers, or other additives. Only use water as the spray carrier.	0
Brassica: Broccoli Brussels Cabbage Cauliflower Kale Rape	Alternaria leaf spot (<i>Alternaria</i> spp.) Black leg (<i>Phoma lingam</i>) Cercospora leaf spot (<i>Cercospora</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.) Ring spot (<i>Mycosphaerella</i> spp.) White leaf spot (<i>Pseudocercospora</i> spp.) Anthracoese (<i>Colletotrichum</i> spp.) White rust (<i>Albugo</i> spp.) Downy mildew (<i>Peronospora</i> spp.) Gray mold (<i>Botrytis</i> spp.) Sclerotinia stem rot (<i>Sclerotinia</i> spp.) Rhizoctonia bottom rot (<i>Rhizoctonia solani</i>)	800 ml	2	Begin applications of Prestige 38 SC prior to disease development and continue on a 7-day to 14-day interval. Use the shorter interval when disease pressure is high. To limit the potential for development of resistance, DO NOT make more than two sequential applications of Prestige 38 SC before alternating to a registered fungicide with a different mode of action. Minimum time from application to harvest (PHI) for leafy brassicas such as Kale and Rape is 14 days.	0 (14)
Bulb / Root Vegetables: Beet root Garlic Ginger Leeks Onions Shallots Turnip	Botrytis leaf blight (<i>Botrytis</i> spp.) Botrytis neck rot (<i>Botrytis</i> spp.) Purple blotch (<i>Alternaria porri</i>) Leaf blight (<i>Alternaria porri</i>) Stemphylium leaf blight (<i>Stemphylium</i> spp.) Downy mildew (<i>Peronospora destructor</i>) Stalk rot (<i>Stemphylium vesicarium</i>)	300 – 600 ml	6	Begin applications prior to onset of disease and continue on a 14-day interval. If application intervals shorter than 14 days are needed, rotate to another fungicide with a different mode of action. Use the higher rate when disease pressure is high. If downy mildew pressure is high, follow the Prestige 38 SC application with a downy mildew fungicide with a different mode of action.	7
Carrots	Alternaria leaf spot (<i>Alternaria</i> spp.) Cercospora leaf spot (<i>Cercospora</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.) Suppression Only: Root rot (<i>Sclerotium rolfsii</i>)	300 – 600 ml	6	Commence applications prior to onset of disease development and continue on a 7-day to 14-day interval. Use the higher rate and the shorter interval when disease pressure is high. To manage resistance, DO NOT make more than two sequential applications of Prestige 38 SC. Alternate with a registered fungicide with a different mode of action. No restriction on livestock grazing or feeding for carrot culls.	0
Citrus: Grapefruit Lemon Lime Mandarin Orange	Alternaria brown spot (<i>Alternaria</i> spp.) Citrus black spot (<i>Guignardia citricarpa</i>) Greasy spot (<i>Mycosphaerella citri</i>) Melanose (<i>Diaporthe citri</i>) Scab (<i>Elsinoe fawcetti</i>)	650 – 800 ml	4	Apply Prestige 38 SC in a regularly scheduled protective fungicide program. Begin applications prior to infection and continue on a 10-day to 21-day interval. Use the higher rate and shorter interval when disease pressure is high. For aerial application to citrus fruit trees, use no less than 90 litres of water per hectare. DO NOT make more than two sequential applications of Prestige 38 SC. Alternate with a registered fungicide with different mode of action.	0
Cotton	Alternaria leaf spot (<i>Alternaria</i> spp.) Anthracoese boll rot (<i>Glomerella</i> spp.) Ascochyta blight (<i>Ascochyta</i> spp.) Ascochyta boll rot (<i>Ascochyta</i> spp.) Cercospora leaf spot (<i>Cercospora</i> spp.) Diplodia boll rot (<i>Diplodia</i> spp.) Hard lock, boll rot (<i>Fusarium</i> spp.)	300 – 800 ml	2	Commence application prior to onset of disease on a 7-day to 14-day interval. Use the higher rate and the shorter interval when disease pressure is high. To limit the potential for development of resistance, DO NOT make more than one application of Prestige 38 SC before alternating to a registered fungicide with a different mode of action.	30

Cucurbits: Butternut Cucumber Pumpkin Squash Watermelon	Alternaria blight (<i>Alternaria cucumerina</i>) Anthracnose (<i>Colletotrichum orbiculare</i>) Cercospora leaf spot (<i>Cercospora citrulina</i>) Downy mildew (<i>Pseudoperonospora</i> spp.) Gummy stem blight (<i>Didymella bryoniae</i>) Powdery mildew (<i>Erysiphe cichoracearum</i> , <i>Sphaerotheca fuliginea</i>)	600 – 800 ml	4	Begin applications prior to onset of disease development and continue on a 7-day to 14-day interval. Use the higher rate and the shorter interval when disease pressure is high. Use the higher rate for anthracnose. To minimize the likelihood of crop injury, it is advisable to do compatibility test and test for crop safety on a small portion of the crop. DO NOT make more than one application of Prestige 38 SC before alternating to a fungicide with a different mode of action.	0
Fruit vegetables: Cherries Chillies Eggplant Okra Paprika Pepper	Black mold (<i>Alternaria alternata</i>) Early blight (<i>Alternaria solani</i>) Late blight (<i>Phytophthora infestans</i>)	300 – 500 ml	6	Begin applications prior to disease development and continue on a 7-day to 14-day intervals. For control of late blight, begin applications prior to disease development; then follow up with a registered fungicide with a different mode of action 5 to 7 days later. Use the higher rate and shorter interval when disease pressure is high.	0
Grapes	Angular leaf spot (<i>Mycosphaerella angulata</i>) Anthracnose (<i>Elsinoe ampelina</i>) Black rot (<i>Guignardia bidwellii</i>) Downy mildew (<i>Plasmopara viticola</i>) Leaf blight (<i>Pseudocercospora vitis</i>) Phomopsis cane and Leaf spot (<i>Phomopsis viticola</i>) Powdery mildew (<i>Uncinula necator</i>) Ripe rot (<i>Colletotrichum gloeosporioides</i>)	600 – 800 ml	5	For <u>powdery mildew</u> control, begin applications as of bud break prior to onset of disease on a 10-day to 14-day interval. Use higher rate on a 14-day to 21-day interval. For <u>black rot and downy mildew</u> control, begin applications as of pre-bloom prior to onset of disease and continue applications on a 10-day to 14-day interval. For control of <u>Botrytis gray mold</u> , apply 800ml/ha prior to onset of disease development when conditions favour disease development during early bloom, bunch pre-closure and veraison up to 14 days before harvest. For all <u>other diseases</u> , begin applications of prior to onset of disease and continue applications on a 10-day to 14-day interval.	14
Herbs /Flowers: Coriander, Hemp Lavender, Lemongrass Marigold, Thyme Ornamentals Parsley, Roses	Phoma blight (<i>Cercosporidium</i> spp.) Powdery mildew (<i>Erysiphe</i> spp., <i>Sphaerotheca</i> spp.) Rust (<i>Puccinia</i> spp.)	600 – 800 ml	3	Begin Prestige 38 SC applications prior to the onset of disease development and repeat application 7 days later as needed or rotate to another fungicide with a different mode of action.	0
Hops	Powdery mildew (<i>Erysiphe cichoracearum</i> , <i>Sphaerotheca</i> spp.) Downy mildew (<i>Pseudoperonospora humuli</i>)	800 ml	3	Begin applications prior to disease development and continue on a 10-day to 21-day interval. Use the shorter interval when disease pressure is high. Adjust water volume to maintain thorough coverage.	14
Leaf greens: Amaranth Blackjack Lettuce Spinach	Alternaria leaf spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Ascochyta leaf spot (<i>Ascochyta</i> spp.) Cercospora leaf spot (<i>Cercospora</i> spp.) Downy mildew (<i>Peronospora</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.) Rust (<i>Puccinia</i> spp.) Septoria leaf spot (<i>Septoria</i> spp.)	300 – 600 ml	2	Begin applications prior to the onset of disease development and continue on a 7-day interval. Use the higher rate when disease pressure is high. To limit the potential for development of resistance, DO NOT make more than one application of Prestige 38 SC before alternating to a registered fungicide with a different mode of action for at least one application.	14
Oil Seeds: Canola Castor oil Sesame Sunflower	Alternaria leaf spot (<i>Alternaria</i> spp.) Cercospora leaf spot (<i>Cercospora helianthini</i>) Downy mildew (<i>Plasmopara halstedii</i>) Powdery mildew (<i>Erysiphe cichoracearum</i>) Rust (<i>Puccinia helianthini</i> , <i>Uromyces</i> spp.) Sclerotinia rot and blight (<i>Sclerotinia</i> spp.) Septoria leaf spot (<i>Septoria</i> spp.)	600 – 800 ml	2	For optimal disease control, begin applications prior to disease development and continue on a 7-day to 14-day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. A second application may be made if disease persists or if weather conditions are favourable for disease development.	21
Peas / Pulses: Cowpeas Fine beans Green beans Groundnuts Lupin Mange tout peas Sugar beans	Alternaria leaf spot (<i>Alternaria</i> spp.) Alternaria pod spot (<i>Alternaria</i> spp.) Ascochyta blight (<i>Ascochyta</i> spp.) Cercospora leaf spot (<i>Cercospora</i> spp.) Downy mildew (<i>Phytophthora nicotianae</i>) Mycosphaerella blight (<i>Mycosphaerella</i> spp.) Powdery mildew (<i>Erysiphe polygami</i>) Rust (<i>Uromyces appendiculatus</i>)	500 – 800 ml	2	For optimal disease control, begin applications prior to onset of disease development or at the beginning of flowering and repeat on a 5-day to 14-day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. DO NOT make more than one application of Prestige 38 SC before alternating to a registered fungicide with a different mode of action for at least one application.	21

	Septoria leaf spot (<i>Septoria</i> spp.)				
Pome fruit: Apples Pears	Alternaria blotch (<i>Alternaria mali</i>) Apple scab (<i>Venturia inaequalis</i>) Bitter root (<i>Colletotrichum</i> spp.) Black rot (<i>Botryosphaeria obtuse</i>) Blue mold (<i>Penicillium</i> spp.) Gray mold (<i>Botrytis</i> spp.) Pear scab (<i>Venturia pirina</i>) Powdery mildew (<i>Podosphaera leucotricha</i>) Sooty blotch (disease complex) White rot (<i>Botryosphaeria dothidea</i>)	600 – 800 ml	4	For <u>scab, powdery mildew, frogeve leaf spot and rust</u> , begin applications prior to disease development and continue on a 7-day to 10-day interval. For <u>blue mold, gray mold, sooty blotch, white rot, black rot, bitter rot and Alternaria blotch</u> , begin applications prior to disease development and continue on a 7-day to 14-day interval. Use the higher rate and shorter interval. For aerial application to pome fruit trees, use no less than 90 litres of spray solution per hectare. DO NOT make more than two sequential applications of Prestige 38 SC before alternating to a registered fungicide with a different mode of action.	0
Potato	Anthracnose (<i>Colletotrichum</i> spp.) Black mold (<i>Alternaria alternata</i>) Botrytis gray mold (<i>Botrytis cinerea</i>) Early blight (<i>Alternaria solani</i>) Late blight (<i>Phytophthora infestans</i>) Powdery mildew (<i>Leveillula Taurica</i>)	300 – 500 ml	3	Begin applications prior to disease development and continue on a 7-day to 14-day interval for anthracnose, black mold, botrytis gray mold and powdery mildew. For control of late blight, begin applications prior to disease development; then follow up with a registered fungicide with a different mode of action 5 to 7 days later. Use the higher rate and shorter interval when disease pressure is high.	7
Stone fruit: Apricot Nectarine Peaches Plums	Alternaria leaf spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Blossom blight (<i>Monilinia</i> spp.) Brown rot (<i>Monilinia</i> spp.) Leaf spot (<i>Blumeriella jaapii</i>) Powdery mildew (<i>Podosphaera</i> spp., <i>Sphaerotheca</i> spp.) Ripe fruit rot (<i>Botrytis cinerea</i> , <i>Monilinia fructicola</i> , <i>Monilinia laxa</i> , <i>Rhizopus</i> spp.) Rust (<i>Tranzschelia discolor</i>) Scab (<i>Cladosporium carpophilum</i>) Shothole (<i>Wilsonomyces carpophilus</i>)	600 – 800 ml	5	Begin application of Prestige 38 SC at pink bud or prior to onset of disease development and continue on a 7-day to 14-day interval. Use the shorter interval and/or the higher rate when disease pressure is high. For aerial application to stone fruit trees, use no less than 90 litres of spray solution per hectare. To manage the potential for development of resistance, DO NOT make more than two sequential applications of Prestige 38 SC or other Group 7 or Group 11 fungicides before alternating to a registered fungicide with a different mode of action.	0
Tomato	Anthracnose (<i>Colletotrichum</i> spp.) Black mold (<i>Alternaria alternata</i>) Botrytis gray mold (<i>Botrytis cinerea</i>) Early blight (<i>Alternaria solani</i>) Late blight (<i>Phytophthora infestans</i>) Powdery mildew (<i>Leveillula Taurica</i>) Septoria leaf spot (<i>Septoria lycopersici</i>) Target spot (<i>Corynespora cassicola</i>)	300 – 500 ml	4	Begin applications prior to disease development and continue on a 7-day to 14-day interval for anthracnose, black mold, botrytis gray mold, early blight, powdery mildew, Septoria leaf spot, and target spot. For control of late blight, begin applications prior to disease development; then follow up with a registered fungicide with a different mode of action 5 to 7 days later. Use the higher rate and shorter interval when disease pressure is high.	
Tree nuts: Almond Cashew nuts Macadamia nuts Pecan nuts Walnuts	Alternaria leaf spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp., <i>Marssonina juglandis</i>) Blossom blight (<i>Monilinia</i> spp.) Botrytis blossom (<i>Botrytis cinerea</i>) Shoot blight (<i>Botrytis cinerea</i>) Green fruit rot (<i>Botrytis cinerea</i>) Leaf rust (<i>Tranzschelia discolor</i>) Panicle / shoot blight (<i>Botryosphaeria</i> spp.) Scab (<i>Cladosporium</i> spp.) Husk spot (<i>Pseudocercospora macadamiae</i>)	800 ml	4	For almonds, begin applications at pink bud and continue on a 7-day to 14-day interval up to 25 days before harvest. For Macadamia, begin application as preventative spray at flowering at 7-dat to 14-day intervals up to a maximum of three applications. In pecan, begin applications prior to onset of disease development and continue on a 7-day to 21-day interval for the control of scab. For all other crops listed, apply Prestige 38 SC prior to disease development and continue on a 7-day to 28-day interval. In all cases, use the shorter interval when shoot growth is very rapid. Use the shorter interval and/or the higher rate when disease pressure is high. PHI for Almond is 25 days.	14 (25)
Tropical fruit: Avocado Banana Dates Guava Kiwi Mango Papaya Pineapple	Anthracnose (<i>Colletotrichum Gloeosporioides</i>) Black spot (<i>Alternaria alternata</i> , <i>Asperisporium caricae</i> , <i>Cercospora papaya</i>) Dry rot (<i>Mycosphaerella</i> spp.) Powdery mildew (<i>Erysiphe</i> spp., <i>Oidium</i> spp.) Pseudocercospora spot (<i>Cercospora</i> spp.) Pseudocercospora blotch (<i>Pseudocercospora Purpurea</i>) Scab (<i>Elsinoe mangiferae</i>)	800 ml	2	Begin application of Prestige 38 SC prior to the onset of disease development and repeat application 7 days later, as needed, or alternate with another registered fungicide having a different mode of action. DO NOT make more than two sequential applications of Prestige 38 SC or other Group 7 or Group 11 fungicides before alternating to a fungicide with a different mode of action. DO NOT apply Prestige 38 SC as a tank mix with any pesticide, fertilizer, nutrient or additive other than non-ionic surfactant.	0