

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE
PRECAUTIONS

1. Handle with care. Avoid splashing and contact with eyes, nose and mouth.
2. Poisonous by swallowing, inhalation and contact with skin.
3. **WEAR FULL PERSONAL PROTECTIVE EQUIPMENT (PPE)** i.e. rubber gloves, rubber boots, overalls, hood, goggles and respirator when preparing and using with this product.
4. **KEEP APART FROM FOOD AND FOOD STUFFS.**
5. **DO NOT** eat, drink or smoke while preparing and applying this product.
6. On completion of spraying remove personal protective clothing, wash face, hands and body with soap and running water before engaging in any activities.
7. **KEEP OUT OF REACH OF CHILDREN AND UNIFORMED PERSONS.**
8. Dangerous to livestock, domestic pets, birds, fish and bees.
9. Store in the original container and **KEEP UNDER LOCK AND KEY.**
10. **DO NOT** contaminate drinking pools, streams, water bodies with the chemical or empty containers.
11. Prevent drift onto other crops and areas not under treatment.

SYMPTOMS OF POISONING:

Nausea, abdominal pain, vomiting, headache, agitation, seizures, hypotension, hypertension, dyspnea, tingling and pruritus on the face and other exposed skin areas.

FIRST AID:

Inhalation: Move victim to fresh air. If not breathing properly, give artificial respiration. Seek medical attention.

Skin contact: Remove contaminated clothing and shoes and launder. Wash thoroughly with soap and running water.

Eye contact: Flush immediately with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

Ingestion: If swallowed, wash the mouth with water provided person is conscious. Induce vomiting after giving a glass of water. Obtain medical attention immediately.

Call Drug and Toxicology Information Services (DATIS) on +263242307148/+26378610098 for more information.

NOTE TO PHYSICIAN:

Antidote: There is no specific antidote. Treat symptomatically.

Rainprita 300 OD

Reg. No.:
23-D-121-14-D-142-121



CONTENTS:
100ML

Composition:.....	mass/volume
Beta-cyfluthrin:.....	90 g/l
Imidacloprid:.....	210 g/l
Inert ingredients:.....	700 g/l

Chemical group:

Beta-cyfluthrin:.....	Synthetic Pyrethroid
Imidacloprid:.....	Neonicotinoid

A broad – spectrum insecticide with systemic, translaminar and contact action for the control of Aphids, Thrips, Whiteflies, Leaf miners and Lepidoptera pests on a wide range of crops.

Shelf Life: Two years from manufactured date if stored in a cool dry place in original unopened container.

Date of Manufacture:

Expiry Date:

Batch No.:



EMPTY CONTAINER DISPOSAL

Invert the container over the spray tank or mixing tank. Allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water to a minimum of 25% of the container. Add the rinsate to the contents of the spray tank. Destroy the empty container by perforation and flattening. **DO NOT** use for any other purposes. If possible, offer the container for recycling and reuse.

CLEANING OF APPLICATION EQUIPMENT

After use, ensure traces of Rainprita 300 OD are removed. Use the following method:

- a) Drain and rinse tank, spray boom and hoses with clean water for at least 10 minutes [rinse at least 3 times].
- b) Fill tank with clean water and add to it 1 liter of household bleach (Sodium hydrochlorite,5%), or 1.5 liter household bleach (sodium hypochlorite 3.5%) per 200 liters of water. Rinse hoses and spray boom and leave in tank for 15minutes whilst agitating. Drain through nozzle outlets.
- c) Repeat step (b) and after that, rinse thoroughly with water. Dispose of the wash water safely in a collection sump where it will not contaminate crops, grazing, rivers, dams and boreholes.

WARRANTY:

The excellent results which this chemical normally yields are not however warranted or guaranteed as the effectiveness can be greatly influenced by factors outside the control of either the manufacturer or the distributor. No warranty whatsoever expressed or implied is therefore given concerning the effectiveness of this chemical. Any responsibility is specifically excluded for any damage, injury or loss of any kind of loss during or resulting from its handling, use or storage whether or not such has been in accordance with label directions.

Manufactured by:

Shandong Weifang Rainbow Chemical Co., Ltd.
Binhai Economic Development Area, Weifang, Shandong, China
Tel: +86-531-88875222, 88875231

Distributed By:

Trowbar Enterprises (PYT) LTD. T/A MAGUIRES
196 Harare Drive, Borrowdale, Harare, Zimbabwe
+263 (0242) 852 131



DIRECTION FOR USE: USE AS DIRECTED.

MIXING INSTRUCTIONS

Half fill the spray container with clean water. Add the required amount of Rainprita 300 OD to the spray container and top up with water to the required volume while agitating continuously. Use the prepared mixture immediately. **DO NOT** prepare more spray mixture than is needed for immediate application. **DO NOT** allow to stand overnight.

WARNING

Avoid repeated use of an insecticide or insecticides from the same chemical group. Alternate or tank-mix with products with different modes of action.

APPLICATION RATES:

Crop	Pest	Dosage/ha in 200L of water/ha	Application directions
Dry Beans	American bollworm, Aphids, Russian Aphids, Leafminers (<i>Liriomyza</i>), Thrips	170-250 mL/ha 16L knapsack sprayer: 14-20 mL	For control of the insects listed below, apply the specified amount with a sprayer in sufficient water to achieve thorough coverage of the attacked crop. Apply when the insects first appear and repeat as
Wheat	Aphids, Colorado Beetle, Leafminers (<i>Liriomyza</i>), caterpillars, armyworm, bollworm		
Tomatoes, Potatoes	Aphids, Whiteflies, Leafminers (<i>Liriomyza</i> and <i>Tuta Absoluta</i>), Thrips	250 mL/ha 16L knapsack sprayer: 20 mL	

Crop	Pest	Dosage/ha in 200L of water/ha	Application directions
Cotton	Aphids, Russian Aphids, Thrips, Mirids, Leafminer (<i>Liriomyza</i>)	170-250 mL/ha 16L knapsack sprayer: 14-20 mL	infestations warrant. DO NOT apply when crops have reached harvest time.
Chili	Girdle beetle and semilooper		
Soybeans	Jassids and Whitefly		

PRE-HARVEST INTERVAL (PHI):

Beans and tomatoes.....14 days
Cotton.....14 days

RE-ENTRY PERIOD

DO NOT enter the treated area for at least 12 hours after treatment.

COMPATIBILITY

Compatible with most of the commonly used pesticides and other crop protection products. However, all mixtures should be tested prior to field application.