

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE

#### WARNINGS

1. Do not sow CUCURBITS in soils treated with **Ronstar 250 EC** for the first six months after application.
2. Avoid spray on to adjacent crops.
3. Do not leave prepared sprays standing overnight.

#### COMPATIBILITY

**Ronstar 250 EC** can safely be tank mix with **Lasso EC**

#### PRECAUTIONS

1. Handle with care. Avoid splashing and contact with the skin.
2. Do not eat, drink or smoke while applying this herbicide.
3. On completion of spraying, wash hands and face thoroughly with soap and water.
4. Maybe dangerous to WILD ANIMALS, BIRDS, FISH, BEES AND OTHER ANIMALS.
5. DO NOT CONTAMINATE DRINKING POOLS, DAMS, RIVERS AND WATERWAYS AND OTHER WATER SUPPLIES.
6. KEEP OUT OF REACH OF CHILDREN.
7. KEEP APART FROM FOOD AND FOODSTUFFS.
8. Store in original container and "KEEP UNDER LOCK AND KEY"
9. Destroy empty container after triple rinsing by burning or burying in a suitable location - do not use for any other purpose.
10. Keep away from heat and open flame.
11. Clean applicator thoroughly after use.
12. Dispose of wash water in a safe manner.
13. **RE-ENTRY** - Do not enter treated area until spray deposit has dried unless wearing protective clothing.

#### SYMPTOMS OF POISONING

No known symptoms.

#### FIRST AID

After skin contact, remove contaminated clothing and wash skin thoroughly with soap and water.

In case of eye contact, irrigate for 20 minutes and seek medical advice.

Should accidental poisoning occur, contact a doctor immediately and show him this label.

#### ADVICE TO PHYSICIAN

Do not induce vomiting. Administer symptomatic and supportive treatment.

#### WARRANTY

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the weeds to the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

#### Packed For:

Bayer Zimbabwe (Pvt) Ltd  
P.O. Box AY 78, Amby, Harare, Zimbabwe.  
Tel. 487211/487245/7 Fax. 487242

Emergency Tel. +27(21) 931-6129

Manufactured by Bayer CropScience, Germany / France

**Ronstar®** is a registered trademark of Bayer CropScience AG, Germany

Date of Manufacture : See on pack

Batch No.: See on pack



Bayer CropScience

Zim/1109/Ronstar 250 EC 5L/Full text label  
Code: 05335832B



Reg. No. 01-C-101-3



POISON

#### COMPOSITION :

	mass/volume
oxadiazon .....	250 g/l
inert ingredients .....	750 g/l

Chemical Group : oxadiazole

A SELECTIVE HERBICIDE FOR THE CONTROL OF  
BROADLEAF WEEDS AND CERTAIN  
ANNUAL GRASSES



CAUTION



**DIRECTIONS FOR USE:** USE ONLY AS DIRECTED

Fill the spray tank with the required quantity of water and the recommended quantity of **Ronstar 250 EC**.

Ensure thorough mixing before spraying commences.

**Ronstar 250 EC** should be applied to a moist soil surface in good tilth before emergence of the weeds.

Where **Ronstar 250 EC** is applied to a dry soil surface, irrigation or rainfall is required immediately after application to activate the herbicide. Regular irrigation or rainfall which will keep the upper soil layer moist will ensure that **Ronstar 250 EC** is kept activated in the soil. After application of **Ronstar 250 EC** the soil surface should be disturbed as little as possible as this will affect the residual activity of the herbicide.

**Ronstar 250 EC** should be applied in a minimum of 300 litres of water per hectare to ensure effective and even coverage. A **Ronstar 250 EC/Lasso** tank-mix for use in soyabeans can be applied by air in a minimum of 40 litres of water per hectare.

CROP	RONSTAR 250 EC APPLICATION RATE	DIRECTIONS
COFFEE	4-5 ℓ / ha	Use the lower rate on soils with less than 20% clay. <b>Ronstar 250 EC</b> maybe applied over the top of coffee seedlings after transplanting. However, weeds emerging close to the base of the stem will be protected by the canopy and thus not controlled. For this reason it is recommended that a directed spray is applied at the base of the coffee seedling by knapsack sprayer.
ONIONS	<b>Direct Seeded:</b> 4-5 ℓ / ha	Apply only after emergence of the onion seedlings and not before the 3-4 leaf stage.
	<b>Sett Planted:</b> 4-5 ℓ / ha	Can be applied either before or after emergence of the onion setts.
	<b>Transplanted:</b> 4-5 ℓ / ha	Apply after transplanting, preferably while the weather is hot and sunny. Irrigate immediately after application.
LAWNS	2% solution	Spot spray patches of <i>Oxalis</i> in lawns with this solution. The lawn may go brown after application but will soon green up again. Use only on couch-type or Kikuyu lawns.
SOYABEANS	It is recommended that <b>Ronstar 250 EC</b> should be applied in a tank-mix with <b>Lasso</b> at the following rates : Soils with less than 20% clay: Not recommended Soils with more than 20% clay: 3 litres <b>Ronstar 250 EC</b> + 2,5-3,0 litres <b>Lasso</b> per hectare.	Apply combination to moist soil after planting but before crop emergence.  Use a higher rate of Lasso where <i>Commelina benghalensis</i> (Wandering Jew) is a problem.

**N.B.** Under certain conditions **Ronstar 250 EC** can cause phytotoxicity of soyabean seedlings. This phytotoxicity takes the form of crinkled leaves and stunting and can vary in degree from mild to severe. However, where this damage occurs, it is only of a transitory nature. The soyabeans grow out of the damage symptoms very quickly and it is unlikely that any yield loss will occur.

CROP	RONSTAR 250 EC APPLICATION RATE	DIRECTIONS
TEA	4-5 ℓ / ha	Use the lower rate in soils with less than 20% clay. <b>Ronstar 250 EC</b> maybe applied over the top of tea seedlings immediately after transplanting.
CARNATIONS	6 ℓ / ha	Apply after transplanting of rooted carnation seedlings. <b>Ronstar 250 EC</b> maybe applied to carnations throughout the growing season
ROSES	4-5 ℓ / ha	Apply to soil after pruning. When spraying is carried out during the growing season, avoid contact of the herbicide with the leaves and young buds. <b>Do not apply to roses grown indoors.</b>
GLADIOLI	4-5 ℓ / ha	Apply after planting of corms but before emergence. Do not apply to corms smaller than 8 cm in circumference

**Lasso® is the trademark of Monsanto Company**

Amongst others the following weeds are normally controlled when **Ronstar 250 EC** is used as directed.

*Amaranthus* spp. - Pigweed

*Chenopodium* spp. - Fat hen

*Commelina benghalensis* - Wandering Jew

*Datura stramonium* - Thorn apple

*Digitaria sanguinalis* - Finger-grass

*Eleusine indica* - Rapoko grass

*Emex australis* - Spiny Emex (Devils thorn)

*Galingsoga parviflora* - Gallant soldier

*Nicandra physaloides* - Apple of Peru

*Oxalis* spp. - Sorrel

*Panicum laevisfolium* - Vlei Panicum

*Portulaca oleracea* - Portulaca

*Richardia brasiliensis* - Mexican clover