


Sunzone Xtra 75 WG

Reg. No.:
23-C-143-28
TRB. No.:
26-26-C-03

CONTENTS:
500G



HARMFUL IF SWALLOWED

Composition:.....mass/volume

Sulfentrazone.....750 g/kg

Inert ingredients: 250 g/kg

Chemical group: Triazolinone

A water dispersible granule herbicide for the control of weeds in crops as indicated.

Manufactured by:

Shandong Weifang Rainbow Chemical Co., Ltd.

Binhai Economic Development Area, Weifang, Shandong, China

Tel: +86-531-88875222, 88875231

Distributed By:

Trowbar Enterprises (PVT) LTD. T/A MAGUIRES

196 Harare Drive, Borrowdale, Harare, Zimbabwe

+263 (0242) 852 131

Date of Manufacture:

Expiry Date:

Batch No.:


Rainbow
all about growing



HERBICIDE

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE

PRECAUTIONS

1. Handle with care. Avoid splashing and contact with eyes, nose and mouth.
2. Poisonous by swallowing, inhalation and contact with skin.
3. **Wear full personal protective equipment (PPE)** i.e. rubber gloves, rubber boots, overalls, hood, goggles and respirator when handle with this product.
4. **KEEP APART FROM FOOD AND FOOD STUFFS.**
5. **DO NOT** eat, drink or smoke while preparing and applying this herbicide.
6. **KEEP OUT OF REACH OF CHILDREN AND UNIFORMED PERSONS.**
7. Dangerous to livestock and domestic pets.
8. Store in the original container and keep under lock and key.
9. Do not contaminate drinking pools, streams, water bodies with the chemical or empty containers.
10. **TOXIC TO FISH AND BIRDS.**
11. Prevent drift onto other crops and areas not under treatment.

SYMPTOMS OF POISONING:

Hazards to humans and domestic animals. Harmful if swallowed, inhaled, or absorbed through skin. Moderate inhalation toxicity. Moderately irritating to the eyes.

Effects from overexposure result from inhaling this product. Symptoms of overexposure include convulsions, tremors, increased sensitivity to touch and sound, labored breathing, decreased locomotion, tearing, nasal discharge and incoordination.

FIRST AID:

Inhalation: Move victim to fresh air. If not breathing properly, give artificial respiration. If breathing is difficult, give oxygen. Consult medical personnel.

Skin contact: Wash thoroughly with soap and running water. Remove contaminated clothing and shoes and launder.

Eye contact: Flush immediately with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

Ingestion: If swallowed, wash the mouth with water provided person is conscious. Induce vomiting. Obtain medical attention immediately.

Call Drug and Toxicology Information Services (DATIS) on +263242307148/+263786100098

NOTE TO PHYSICIAN:

Antidote: There is no specific antidote. Treat symptomatically.

EMPTY CONTAINER DISPOSAL

Invert the container over the spray tank or mixing tank. Allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water to a minimum of 10% of the container. Add the rinsate to the contents of the spray tank. Destroy the empty container by perforation and flattening. Bury away in a designated area or away from water sources. DO NOT use for any other purposes. If possible, offer the container for recycling and reuse.

CLEANING OF APPLICATION EQUIPMENT

After use, ensure traces of Sunzone Xtra 75 WG are removed. Use the following method:

- a) Drain and rinse tank, spray boom and hoses with clean water for at least 10 minutes (rinse at least 3 times).
- b) Fill tank with clean water and add to it 1 liter of household bleach (Sodium hydrochlorite, 5%) per 200 liters of water. Rinse hoses and spray boom and leave in tank for 15 minutes whilst agitating. Drain through nozzle outlets.
- c) Repeat step (b) and after that, rinse thoroughly with water. Dispose of the wash water safely in a collection sump where it will not contaminate crops, grazing, rivers, dams and boreholes.

WARRANTY:

The excellent results which this chemical normally yields are not however warranted or guaranteed as the effectiveness can be greatly influenced by factors outside the control of either the manufacturer or the distributor. No warranty whatsoever expressed or implied is therefore given concerning the effectiveness of this chemical. Any responsibility is specifically excluded for any damage, injury or loss of any kind of loss during or resulting from its handling, use or storage whether or not such has been in accordance with label directions.

DIRECTION FOR USE:

USE AS DIRECTED.

USE RESTRICTIONS

Follow up crops: **Do not** plant any other crop, apart from sugarcane, within 24 months after application of Sunzone Xtra 75 WG. Ensure that the soil is ploughed thoroughly and harrowed before any other crop is planted.

MIXING INSTRUCTIONS

Half fill the spray container with clean water. Add the required amount of Sunzone Xtra 75 WG to the spray container and top up with water to the required volume while agitating continuously.

Use the prepared mixture immediately. Do not prepare more spray mixture than is needed for immediate application. Do not allow to stand overnight.

APPLICATION RATES:

Crop	Dosage/ha	Application directions
Tobacco	0-20% Clay: 270-300g >20% Clay: 330-350g	Yellow and purple nutsedge, and B/L weeds. Use higher rates for improved B/L weed control. Apply within 4 days of transplanting. Avoid spraying hearts of plants.
Sugarcane (Plant and Ratoon Crops)	1.2kg	Pre-emergence treatment Apply Sunzone Xtra 75 WG on its own. Application must be done onto clean, cultivated soil before emergence of the weeds. Apply onto moist soil, rain or irrigation is required within one week after application to obtain best results. Pre-emergence application of Sunzone Xtra 75 WG may result in slight scorching of the leaves, especially if the crop is subjected to stress conditions (e.g. drought). However, these symptoms are only temporary and will disappear as the crop grows.

LIST OF WEEDS CONTROLLED

<i>Ageratum conyzoides</i>	Invading Ageratum
<i>Amaranthus spp</i>	Pigweed
<i>Bulbostylis hispidula</i>	Slender sedge
<i>Commelina bengalensis</i>	Wandering Jew
<i>Conyza spp.</i>	Fleabane
<i>Cyperus esculentus</i>	Yellow nutsedge
<i>Datura stramonium</i>	Thorn apple
<i>Digitaria sanguinalis</i>	Crab finger grass
<i>Emex australis</i>	Spiny emex
<i>Eleusine indica</i>	Goose grass
<i>Panicum maximum</i>	Buffalo grass
<i>Senecio madagascariensis</i>	Madagascar ragwort
<i>Setaria verticillata</i>	Sticky bristle grass
<i>Solanum nigrum</i>	Black nightshade
<i>Sonchus oleraceus</i>	Sowthistle
<i>Sysimbrium thellungii</i>	Common wild mustard

RE-ENTRY PERIOD

DO NOT enter the treated area for at least 24 hours after treatment.

COMPATIBILITY

Compatible with most of the commonly used pesticides and other crop protection products. However all mixtures should be tested prior to field application.