

15 mm 65 mm 27.5 mm 130 mm 27.5 mm 50 mm 15 mm 15 mm

15 mm

250 mm

220 mm

15 mm

**Preparing mixture:**  
Premix the required amount of Trigard® 75 WP with a small amount of water to form a slurry. Half fill the spray tank with clean water and pour the slurry through a sieve into the tank. Add more water to fill the tank up to the final volume.

**Compatibility:**  
Trigard® 75 WP is compatible with most insecticides, acaricides and fungicides. The compatibility of Trigard® 75 WP with third party products may be affected by the formulation of the other products and / or the quality of the water used for dilution. Since Syngenta cannot be aware of the water quality, we cannot accept responsibility for poor results or crop damage. Test compatibility with the water and the product to be used in the mixture prior to application to ensure both physical compatibility and crop safety.


**Directions for use**  
**Dosage rates**

Crop	Dosage rates	Spray interval and timing
Tomatoes	150 g/ha in 150-1500 litres of water depending on plant height	The first spray should be at the first sign of infestation. Repeat spraying at 7 day intervals if necessary.
Ornamentals	150 g/ha (20 g/100 litres water)	First spray should be at first sign of infestation. Repeat at 7 day intervals if infestation pressure is high and 14 days if it is low.


**Emergency Call Number:**  
Syngenta Alarm Centre in Huddersfield, UK  
Tel.: +44 1484 538 444, Fax: +44 1484 554 093

**Distributor:**

L166628 ZIMB/08Q PPE 4014213



**200 grams**



**Composition** m/m  
 Cyromazine ..... 75%  
 Inert ingredients ..... 25%  
 Chemical Group: Triazine

**BEFORE USING THIS PRODUCT READ AND UNDERSTAND THE ENTIRE LABEL.**

**Shelf Life:** At least 2 years if stored in a cool dry place in original unopened container.

**Wettable powder.**

**Trigard® 75 WP is an insecticide growth regulator for the control of leafminer in tomatoes and ornamentals.**


**Registered by: Syngenta Agro Services**  
 P.O. Box 1088, Harare  
 Suite Za, Julia's Parade  
 Sam Levy's Village  
 Borrowdale, Tel.: 0772 402 477

**Manufacturer:** Syngenta Crop Protection AG, Basle, Switzerland

® Registered Trademark of a Syngenta Group Company

© 2007 Syngenta Crop Protection AG

**TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.**



- Precautions**
1. Handle with care; avoid splashing/contact; poisonous by swallowing, inhalation and contact with the skin.
  2. Wear protective clothing, i.e gloves, overall, rubber boots, hood and respirator.
  3. Do not eat, drink or smoke while applying this product.
  4. Remove protective clothing on completion of spraying and wash hands and face thoroughly with soap and water.
  5. Keep out of reach of children.
  6. Keep apart from food, foodstuffs, seeds and fertilizers.
  7. Store in original container in a cool, dry place and KEEP UNDER LOCK AND KEY.
  8. Use only on the crops for which the product is registered.
  9. Avoid drift onto adjacent crops.
  10. After use, wash sprayer out thoroughly and dispose of washings in a safe place, e.g. on waste ground. Do not contaminate water supplies.
  11. Empty container - triple rinse, puncture and place in secure disposal site; DO NOT use for any other purpose.

**Symptoms of poisoning:**  
No case of human poisoning is on record.

**First aid:**  
If poisoning is suspected, stop work immediately and seek medical advice. In case of contact with skin and eyes, wash immediately with plenty of water. Rinse splashes from eyes with abundant fresh and clean water for at least 10 minutes. If inhaled, move to clean air. If swallowed, give a large quantity of medical charcoal in plenty of water.

**Note to physician:**  
No specific antidote is known. Apply symptomatic therapy.

**Warranty:**  
The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

**Mode of action:**  
Trigard® 75 WP is an insect growth regulator and inhibits the development of dipterous larvae; ingestion of the product by the adult flies also results in a reduction in egg number and egg hatch.

**Warning:**  
The pre-harvest interval in tomatoes is 3 days.